




Alain Matthes

July 15, 2022

Abstract

This document brings together some notes about **tkz-euclide 4**, a tool for creating geometric figures. The two most important Euclidean tools used by early Greeks to construct different geometrical shapes and angles were a compass and a straightedge. My idea is to allow you to follow step by step a construction that would be done by hand (with compass and straightedge) as naturally as possible.

Example I

Book I, proposition I __Euclide__

To construct an equilateral triangle on a given finite straight line.

Explanation

The fourth tutorial of the *PgfManual* is about geometric constructions. *T. Tantau* proposes to get the drawing with its beautiful tool *TikZ*. Here I propose the same construction with *tkz-euclide 4*. The color of the *TikZ* code is green and that of *tkz-euclide 4* is red.

```
\usepackage{tikz}
\usetikzlibrary{calc,intersections,through,backgrounds}

\usepackage{tkz-euclide}
```

How to get the line AB ? To get this line, we use two fixed points.

```
\coordinate [label=left:$A$] (A) at (0,0);
\coordinate [label=right:$B$] (B) at (1.25,0.25);
\draw (A) -- (B);
\tkzDefPoint(0,0){A}
\tkzDefPoint(1.25,0.25){B}
\tkzDrawSegment(A,B)
\tkzLabelPoint[left](A){$A$}
\tkzLabelPoint[right](B){$B$}
```

We want to draw a circle around the points A and B whose radius is given by the length of the line AB.

```
\draw let \p1 = ($ (B) - (A) $),
\n2 = {veclen(\x1,\y1)} in
(A) circle (\n2)
(B) circle (\n2);

\tkzDrawCircles(A,B B,A)
```

The intersection of the circles $\mathcal{D}$ and $\mathcal{E}$

```
draw [name path=A--B] (A) -- (B);
node (D) [name path=D,draw,circle through=(B),label=left:$D$] at (A) {};
node (E) [name path=E,draw,circle through=(A),label=right:$E$] at (B) {};
path [name intersections={of=D and E, by={ [label=above:$C$]C, [label=below:$C'$]C' }}];
draw [name path=C--C',red] (C) -- (C');
path [name intersections={of=A--B and C--C',by=F}];
node [fill=red,inner sep=1pt,label=-45:$F$] at (F) {};


\tkzInterCC(A,B)(B,A) \tkzGetPoints{C}{X}
```

How to draw points :

```
\foreach \point in {A,B,C}
\fill [black,opacity=.5] (\point) circle (2pt);

\tkzDrawPoints[fill=gray,opacity=.5](A,B,C)
```

The complete code with tkz-euclide 4



```
\begin{tikzpicture}[thick,help lines/.style={thin,draw=black!50},old paper]
\tkzDefPoint(0,0){A}
\tkzDefPoint(1.25+rand(),0.25+rand()){B}
\tkzInterCC(A,B)(B,A) \tkzGetPoints{C}{X}

\tkzFillPolygon[triangle](A,B,C)
\tkzDrawSegment[input](A,B)
\tkzDrawSegments[red](A,C B,C)
\tkzDrawCircles[help lines](A,B B,A)

\tkzLabelPoints(A,B)
\tkzLabelCircle[below=12pt](A,B)(180){$\mathcal{D}$}
\tkzLabelCircle[above=12pt](B,A)(180){$\mathcal{E}$}
\tkzLabelPoint[above,red](C){$C$}
\tkzDrawPoints[fill=gray,opacity=.5](A,B,C)

\tkzText[text width=8cm,align=justify](0,-4){%
  \small\textbf{Proposition I}\n
  \emph{To construct an \textcolor{triangle}{equilateral triangle}
  on a given \textcolor{input}{finite straight line}.}
  \n
  \vskip1em
  Let \A\B be the given \textcolor{input}{finite straight line}. \dots
}
\end{tikzpicture}
```

Example II

Book I, Proposition II _Euclid's Elements_

To place a straight line equal to a given straight line with one end at a given point.

Explanation

In the first part, we need to find the midpoint of the straight line AB. With **TikZ** we can use the calc library

```
\coordinate [label=left:$A$] (A) at (0,0);
\coordinate [label=right:$B$] (B) at (1.25,0.25);
\draw (A) -- (B);
\node [fill=red,inner sep=1pt,label=below:$X$] (X) at ($ (A)!.5!(B) $) {};
```

With **tkz-euclide** we have a macro `\tkzDefMidPoint`, we get the point X with `\tkzGetPoint` but we don't need this point to get the next step.

```
\tkzDefPoints{0/0/A,0.75/0.25/B,1/1.5/C}
\tkzDefMidPoint(A,B) \tkzGetPoint{X}
```

Then we need to construct a triangle equilateral. It's easy with **tkz-euclide**. With **TikZ** you need some effort because you need to use the midpoint X to get the point D with trigonometry calculation.

```
\node [fill=red,inner sep=1pt,label=below:$X$] (X) at ($ (A)!.5!(B) $) {};
```

```
\node [fill=red,inner sep=1pt,label=above:$D$] (D) at
($ (X) ! {sin(60)*2} ! 90:(B) $) {};
```

```
\draw (A) -- (D) -- (B);
```

```
\tkzDefTriangle[equilateral](A,B) \tkzGetPoint{D}
```

We can draw the triangle at the end of the picture with

```
\tkzDrawPolygon{A,B,C}
```

We know how to draw the circle $\mathcal{H}$ around B through C and how to place the points E and F

```
\node (H) [label=135:$H$,draw,circle through=(C)] at (B) {};
```

```
\draw (D) -- ($ (D) ! 3.5 ! (B) $) coordinate [label=below:$F$] (F);
```

```
\draw (D) -- ($ (D) ! 2.5 ! (A) $) coordinate [label=below:$E$] (E);
```

```
\tkzDrawCircle(B,C)
\tkzDrawLines[add=0 and 2](D,A D,B)
```

We can place the points E and F at the end of the picture. We don't need them now.

Intersecting a Line and a Circle : here we search the intersection of the circle around B through C and the line DB. The infinite straight line DB intercepts the circle but with **TikZ** we need to extend the lines DB and that can be done using partway calculations. We get the point F and BF or DF intercepts the circle

```
\node (H) [label=135:$H$,draw,circle through=(C)] at (B) {};
```

```
\path let \p1 = ($ (B) - (C) $) in
coordinate [label=left:$G$] (G) at ($ (B) ! vecLen(\x1,\y1) ! (F) $);
```


```
\fill[red,opacity=.5] (G) circle (2pt);
```

Like the intersection of two circles, it's easy to find the intersection of a line and a circle with `tkz-euclide`. We don't need F

```
\tkzInterLC(B,D)(B,C)\tkzGetFirstPoint{G}
```

There are no more difficulties. Here the final code with some simplifications. Nous tracons le cercle $\mathcal{K}$ de centre D et passant par G. Il coupe la droite AD au point L. $AL = BC$.

```
\tkzDrawCircle(D,G)
\tkzInterLC(D,A)(D,G)\tkzGetSecondPoint{L}
```



```

\begin{tikzpicture}[scale=2,old paper] % remove old paper style !
\tkzDefPoint(0,0){A}
\tkzDefPoint(0.75,0.25){B}
\tkzDefPoint(1,1.5){C}
\tkzDefTriangle[equilateral](A,B) \tkzGetPoint{D}
\tkzInterLC[near](D,B)(B,C) \tkzGetSecondPoint{G}
\tkzInterLC[near](D,A)(D,G) \tkzGetFirstPoint{L}
\tkzDrawCircles(B,C D,G)
\tkzDrawLines[add=0 and 2](D,A D,B)
\tkzDrawSegment(A,B)
\tkzDrawSegments[red](A,L B,C)
\tkzDrawPoints[red](D,L,G)
\tkzDrawPoints[fill=gray](A,B,C)
\tkzLabelPoints[left,red](A)
\tkzLabelPoints[below right,red](L)
\tkzLabelCircle[above](B,C)(20){$\mathcal{H}$}
\tkzLabelPoints[above left](D)
\tkzLabelPoints[above](G)
\tkzLabelPoints[above,red](C)
\tkzLabelPoints[right,red](B)
\tkzLabelCircle[below](D,G)(-90){$\mathcal{K}$}
\tkzText[text width=8cm,align=justify](0,-2){%
  \small\textbf{Proposition II}\}
  \emph{To place a straight line equal to a given straight line with one end at a given point.}\}
  Let \A\ be the given point, and \B\C\ the given straight line \dots}
\end{tikzpicture}

```

Example III

Definition of a circle _Apollonius_

To place a straight line equal to a given straight line with one end at a given point.

Explanation

From Wikipedia : *Apollonius showed that a circle can be defined as the set of points in a plane that have a specified ratio of distances to two fixed points, known as foci. This Apollonian circle is the basis of the Apollonius pursuit problem. ... The solutions to this problem are sometimes called the circles of Apollonius.* A circle is the set of points in a plane that are equidistant from a given point O. The distance r from the center is called the radius, and the point O is called the center. It is the simplest definition but it is not the only one. Apollonius of Perga gives another definition : The set of all points whose distances from two fixed points are in a constant ratio is a circle.

With tkz-euclide is easy to show you the last definition

The code and the analyse

```

\documentclass{standalone}
% Excellent class to show the result and to verify the bounding box.


```

```

\usepackage{tkz-euclide}
  % no need to use \usetkzobj !
\begin{document}
\begin{tikzpicture}[scale=1.5]
  % Firstly we defined two fixed point.
  % The figure depends of these points and the ratio K
\tkzDefPoint(0,0){A}
\tkzDefPoint(4,0){B}
  % tkz-euclide.sty knows about the apollonius's circle
  % with K=2 we search some points like I such as  $IA=2 \times IB$ 
\tkzDefCircle[apollonius,K=2](A,B) \tkzGetPoint{K1}
\tkzGetLength{rAp}
\tkzDefPointOnCircle[R = center K1 angle 30 radius \rAp]
\tkzGetPoint{I}
\tkzDefPointOnCircle[R = center K1 angle 280 radius \rAp]
\tkzGetPoint{J}
%\tkzDrawCircle[R,color = teal,fill=MidnightBlue!20,opacity=.4](K1,\rAp pt)
\tkzDrawSegments[red](A,I I,B A,J J,B)
\tkzDrawSegment(A,B)
\tkzLabelPoints[below,font=\scriptsize](A,B,K1,I,J)
\tkzDrawPoints(A,B,K1,I,J)
\end{tikzpicture}
\end{document}

```

The result



Example IV

The Apollonius circle of a triangle *_Apollonius_*

To place a straight line equal to a given straight line with one end at a given point.

Explanation

The purpose of the first examples was to show the simplicity with which we could recreate these propositions. With TikZ you need to do calculations and use trigonometry while with **tkz-euclide** you only need to build simple objects

But don't forget that behind or far above **tkz-euclide** there is TikZ. I'm only creating an interface between TikZ and the user of my package.

The last example is very complex and it is to show you all that we can do with **tkz-euclide**.

The code and the analyse

```
% !TEX TS-program = lualatex
\documentclass{standalone}
\usepackage{tkz-euclide}
\begin{document}

\begin{tikzpicture}[scale=.6,old paper]
\tkzDefPoints{O/O/A,6/O/B,0.8/4/C}
\tkzDefTriangleCenter[euler](A,B,C)\tkzGetPoint{N}
\tkzDefTriangleCenter[circum](A,B,C)\tkzGetPoint{O}
\tkzDefTriangleCenter[lemoine](A,B,C)\tkzGetPoint{K}
\tkzDefTriangleCenter[ortho](A,B,C)\tkzGetPoint{H}
\tkzDefSpcTriangle[excenter,name=J](A,B,C){a,b,c}
\tkzDefSpcTriangle[centroid,name=M](A,B,C){a,b,c}
\tkzDefCircle[in](Ma,Mb,Mc)\tkzGetPoint{Sp} % Sp Spieker center
\tkzDefProjExcenter[name=J](A,B,C)(a,b,c){Y,Z,X}
\tkzDefLine[parallel=through Za](A,B)\tkzGetPoint{Xc}
\tkzInterLL(Za,Xc)(C,B)\tkzGetPoint{C'}
\tkzDefLine[parallel=through Zc](B,C)\tkzGetPoint{Ya}
\tkzInterLL(Zc,Ya)(A,B)\tkzGetPoint{A'}
\tkzDefPointBy[reflection= over Ja--Jc](C')\tkzGetPoint{Ab}
\tkzDefPointBy[reflection= over Ja--Jc](A')\tkzGetPoint{Cb}
\tkzInterLL(K,O)(N,Sp)\tkzGetPoint{Q}
\tkzInterLC(A,B)(Q,Cb)\tkzGetFirstPoint{Ba}
\tkzInterLC(A,C)(Q,Cb)\tkzGetPoints{Ac}{Ca}
\tkzInterLC(B,C')(Q,Cb)\tkzGetFirstPoint{Bc}
\tkzInterLC[next to=Ja](Ja,Q)(Q,Cb)\tkzGetFirstPoint{F'a}
\tkzInterLC[next to=Jc](Jc,Q)(Q,Cb)\tkzGetFirstPoint{F'c}
\tkzInterLC[next to=Jb](Jb,Q)(Q,Cb)\tkzGetFirstPoint{F'b}
\tkzInterLC[common=F'a](Sp,F'a)(Ja,F'a)\tkzGetFirstPoint{Fa}
\tkzInterLC[common=F'b](Sp,F'b)(Jb,F'b)\tkzGetFirstPoint{Fb}
\tkzInterLC[common=F'c](Sp,F'c)(Jc,F'c)\tkzGetFirstPoint{Fc}
\tkzInterLC(Mc,Sp)(Q,Cb)\tkzGetFirstPoint{A''}
```

```

\tkzDefCircle[euler](A,B,C)
\tkzDefCircle[ex](C,A,B)
\tkzDefCircle[ex](A,B,C)
\tkzDefCircle[ex](B,C,A)
\tkzGetPoints{E}{e}
\tkzGetPoints{Ea}{a}
\tkzGetPoints{Eb}{b}
\tkzGetPoints{Ec}{c}
% Calculations are done, now you can draw, mark and label
\tkzDrawCircles(Q,Cb E,e)%
\tkzDrawCircles(Eb,b Ea,a Ec,c)
\tkzDrawPolygon(A,B,C)
\tkzDrawSegments[dashed](A,A' C,C' A',Zc Za,C' B,Cb B,Ab A,Ca)
\tkzDrawSegments[dashed](C,Ac Ja,Xa Jb,Yb Jc,Zc)
\begin{scope}
\tkzClipCircle(Q,Cb) % We limit the drawing of the lines
\tkzDrawLine[add=5 and 12,orange](K,O)
\tkzDrawLine[add=12 and 28,red!50!black](N,Sp)
\end{scope}
\tkzDrawPoints(A,B,C,K,Ja,Jb,Jc,Q,N,O,Sp,Mc,Xa,Xb,Yb,Yc,Za,Zc)
\tkzDrawPoints(A',C',A'',Ab,Cb,Bc,Ca,Ac,Ba,Fa,Fb,Fc,F'a,F'b,F'c)
\tkzLabelPoints(Ja,Jb,Jc,Q,Xa,Xb,Za,Zc,Ab,Cb,Bc,Ca,Ac,Ba,F'b)
\tkzLabelPoints[above](O,K,F'a,Fa,A'')
\tkzLabelPoints[below](B,F'c,Yc,N,Sp,Fc,Mc)
\tkzLabelPoints[left](A',C',Fb)
\tkzLabelPoints[right](C)
\tkzLabelPoints[below right](A)
\tkzLabelPoints[above right](Yb)
\tkzDrawSegments(Fc,F'c Fb,F'b Fa,F'a)
\tkzDrawSegments[color=green!50!black](Mc,N Mc,A'' A'',Q)
\tkzDrawSegments[color=red,dashed](Ac,Ab Ca,Cb Ba,Bc Ja,Jc A',Cb C',Ab)
\tkzDrawSegments[color=red](Cb,Ab Bc,Ac Ba,Ca A',C')
\tkzMarkSegments[color=red,mark=|](Cb,Ab Bc,Ac Ba,Ca)
\tkzMarkRightAngles(Jc,Zc,A Ja,Xa,B Jb,Yb,C)
\tkzDrawSegments[green,dashed](A,F'a B,F'b C,F'c)
\end{tikzpicture}
\end{document}

```

The result

